

> d his full

(FILE 'HOME' ENTERED AT 17:03:47 ON 10 FEB 2006)

FILE 'MEDLINE, AGRICOLA, CABA, CAPLUS, BIOSIS, DISSABS, EMBASE' ENTERED
AT 17:05:05 ON 10 FEB 2006

L1 245 SEA ABB=ON PLU=ON 4G AND 5G AND FIBRINOLYSIS
L2 0 SEA ABB=ON PLU=ON 4G/5G AND FIBRINOLYSIS
L3 274 SEA ABB=ON PLU=ON 4G AND FIBRINOLYSIS
L4 122 SEA ABB=ON PLU=ON L3 AND (EXERCISE OR PHYSICAL OR ACTIVITY)

L5 131 DUP REM L3 (143 DUPLICATES REMOVED)
L6 59 DUP REM L4 (63 DUPLICATES REMOVED)
L7 39 SEA ABB=ON PLU=ON L6 AND PY<=2002
D L7 1- TI
L8 4 SEA ABB=ON PLU=ON L7 AND EXERCISE
D L8 1- TI
D L8 4 IBIB ABS KWIC
E HAGBERG J/AU 21
L9 8 SEA ABB=ON PLU=ON ("HAGBERG J"/AU OR "HAGBERG J M"/AU OR
"HAGBERG JAMES M"/AU OR "HAGBERG JAMES MICHAEL"/AU) AND
FIBRINOLYSIS
D L9 1- TI
L10 1 SEA ABB=ON PLU=ON L9 AND PY<=2002
D L10 1- TI
D L10 1- IBIB
E KULAPUTANA O/AU 21
L11 9 SEA ABB=ON PLU=ON ("KULAPUTANA O"/AU OR "KULAPUTANA ONANONG"/
AU OR "KULAPUTANA ORNANONG"/AU) AND FIBRINOLYSIS
D L11 1I TI
D L11 1- TI
E FERRELL R/AU 21
E FERRELL ROBERT/AU 21



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
Welcome kapus

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search PubMed for 4g/5g fibrinolysis exercise

Preview

Go

C

Limits

Preview/Index

History

Clipboard

Details

Query #1 deleted

- Search History will be lost after eight hours of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.
- Click on query # to add to strategy

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Search

Most Recent Queries

Time Result

#34 Search 4g/5g fibrinolysis exercise	17:00:00	3
#32 Related Articles for PubMed (Select 10494774)	16:59:38	172
#13 Search 4g/5g exercise	16:49:53	4
#31 Search 4g/5g and exercise	16:49:50	4
#30 Search ferrell and exercise	16:49:34	44
#29 Search kulaputana and exercise	16:44:04	6
#28 Search hagberg j and exercise	16:43:47	148
#27 Search hagberg j and genotype	16:43:25	25
#26 Search hagberg j and pai	16:43:20	1
#25 Search hagberg j and 4g	16:43:17	0
#24 Search hagberg j and plasminogen	16:43:03	1
#23 Search hagberg j and plasminogen Limits: Publication Date from 1980 to 2002/07/05	16:42:57	0
#22 Search hagberg j Limits: Publication Date from 1980 to 2002/07/05	16:42:46	127
#21 Search 4g/5g fibrinolysis exercise Limits: Publication Date from 1980 to 2002/07/05	16:40:59	3
#19 Search 4g/5g fibrinolysis Field: All Fields, Limits: Publication Date from 1980 to 2002/07/05	16:39:45	37
#18 Search 4g/5g fibrinolysis	16:17:22	66
#12 Search 4g/5g exercise	15:54:35	0
#11 Search 4g/5g	15:48:31	302

Clear History

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Feb 6 2006 04:30:54

SEARCH	DIGITAL DISSERTATIONS	ProQuest
BROWSE		
RESULTS LIST MARKED LIST SHOPPING CART DOWNLOAD CONTACT		

Full Citation & Abstract

PRICING ?

PUBLICATION NUMBER AAT 3035829

TITLE Effects of a common genetic polymorphism and exercise training on fibrinolysis in men and women aged 50--70

AUTHOR Tiyasangthong, Onanong

DEGREE PhD

SCHOOL UNIVERSITY OF MARYLAND, COLLEGE PARK

DATE 2001

PAGES 172

ADVISER Hagberg, James M.

ISBN 0-493-48931-2

SOURCE DAI-B 62/12, p. 5635, Jun 2002

SUBJECT HEALTH SCIENCES, MEDICINE AND SURGERY (0564); HEALTH SCIENCES, PUBLIC HEALTH (0573); HEALTH SCIENCES, REHABILITATION AND THERAPY (0382)

DIGITAL FORMATS 6.15Mb image-only PDF

ORDER NOW ►

FREE DOWNLOAD

24 page Preview



Prospective epidemiological investigations suggest that impaired fibrinolysis is an independent risk factor for cardiovascular disease. Endurance exercise training has an inconsistent effect on fibrinolysis, possibly due to inadequate study design and lack of control of critical variables. The purpose of this study was to investigate the effects of endurance exercise training on blood fibrinolysis in middle-aged to older men and women. The plasminogen activator inhibitor-1 (PAI-1) gene promoter 4G/5G polymorphism appears to affect circulating PAI-1 levels. The present study also determined if the PAI-1 polymorphism and initial PAI-1 levels play roles in exercise training-induced fibrinolysis changes. Sedentary healthy volunteers aged 50--70 years were studied. Subjects completed a 6-week dietary program before baseline testing. Baseline tests in 48 subjects included fibrinolytic enzymes (tissue plasminogen activator (t-PA) and PAI-1 activities, and t-PA antigen), lipid-lipoprotein profiles, body composition, and maximal oxygen consumption ($\text{VO}_{2\text{max}}$). Genotyping was performed using PCR/RFLP techniques. Baseline tests were repeated in 34 subjects after 6-month exercise intervention. Baseline fibrinolytic levels were not significantly different among genotype groups (15.12 ± 1.39 vs. 11.97 ± 0.73 vs. 11.89 ± 1.39 AU/mL, for 4G/4G, 4G/5G, and 5G/5G, respectively). However, after controlling for triglyceride levels, the 4G/4G genotype had significantly higher PAI-1 activity levels than the other genotypes. After exercise training, fibrinolysis improved as shown by a 30% increase in t-PA activity (0.18 ± 0.07 IU/mL, $p = 0.016$) and a 6% decrease in t-PA antigen (-0.58 ± 0.20 , $p = 0.006$). Subgroup analyses revealed a significant 41% increase ($p = 0.03$) in t-PA activity levels in the 4G/5G group and a 10% decrease in t-PA antigen levels ($p = 0.008$) in the 4G/4G group. Training did not change PAI-1 activity levels significantly. ANOVAs revealed

no differences in fibrinolytic measure changes with training among genotype groups. Adjusting for initial levels did not change results for PAI-1 activity and t-PA antigen. Interestingly, there was a significant interaction between baseline t-PA activity and genotype in determining final t-PA activity. The present results suggest that endurance exercise training may increase fibrinolysis in middle-aged to older men and women by increasing t-PA activity and decreasing t-PA antigen levels. Moreover, individuals with 4G alleles may respond more favorably to training in terms of improving fibrinolysis.

You are viewing title 1 of 2 in the current results list.

NEXT



GO TO NUMBER:

=> d 18 4 ibib abs kwic

L8 ANSWER 4 OF 4 DISSABS COPYRIGHT (C) 2006 ProQuest Information and Learning Company; All Rights Reserved on STN
ACCESSION NUMBER: 2002:35148 DISSABS Order Number: AAI3035829
TITLE: Effects of a common genetic polymorphism and exercise training on fibrinolysis in men and women aged 50--70
AUTHOR: Tiyaangthong, Onanong [Ph.D.]; Hagberg, James M. [adviser]
CORPORATE SOURCE: University of Maryland College Park (0117)
SOURCE: Dissertation Abstracts International, (2001) Vol. 62, No. 12B, p. 5635. Order No.: AAI3035829. 172 pages. ISBN: 0-493-48931-2.
DOCUMENT TYPE: Dissertation
FILE SEGMENT: DAI
LANGUAGE: English

AB Prospective epidemiological investigations suggest that impaired fibrinolysis is an independent risk factor for cardiovascular disease. Endurance exercise training has an inconsistent effect on fibrinolysis, possibly due to inadequate study design and lack of control of critical variables. The purpose of this study was to investigate the effects of endurance exercise training on blood fibrinolysis in middle-aged to older men and women.

The plasminogen activator inhibitor-1 (PAI-1) gene promoter 4G/5G polymorphism appears to affect circulating PAI-1 levels. The present study also determined if the PAI-1 polymorphism and initial PAI-1 levels play roles in exercise training-induced fibrinolysis changes.

Sedentary healthy volunteers aged 50-70 years were studied. Subjects completed a 6-week dietary program before baseline testing. Baseline tests in 48 subjects included fibrinolytic enzymes (tissue plasminogen activator (t-PA) and PAI-1 activities, and t-PA antigen), lipid-lipoprotein profiles, body composition, and maximal oxygen consumption (VO2 max). Genotyping was performed using PCR/RFLP techniques. Baseline tests were repeated in 34 subjects after 6-month exercise intervention.

Baseline fibrinolytic levels were not significantly different among genotype groups (15.12 ± 1.39 vs. 11.97 ± 0.73 vs. 11.89 ± 1.39 AU/mL, for 4G/4G, 4G/5G, and 5G/5G, respectively). However, after controlling for triglyceride levels, the 4G/4G genotype had significantly higher PAI-1 activity levels than the other genotypes. After exercise training, fibrinolysis improved as shown by a 30% increase in t-PA activity (0.18 ± 0.07 IU/mL, $p = 0.016$) and a 6% decrease in t-PA antigen (-0.58 ± 0.20 , $p = 0.006$). Subgroup analyses revealed a significant 41% increase ($p = 0.03$) in t-PA activity levels in the 4G/5G group and a 10% decrease in t-PA antigen levels ($p = 0.008$) in the 4G/4G group. Training did not change PAI-1 activity levels significantly. ANOVAs revealed no differences in fibrinolytic measure changes with training among genotype groups. Adjusting for initial levels did not change results for PAI-1 activity and t-PA antigen. Interestingly, there was a significant interaction between baseline t-PA activity and genotype in determining final t-PA activity.

The present results suggest that endurance exercise training may increase fibrinolysis in middle-aged to older men and women by increasing t-PA activity and decreasing t-PA antigen levels. Moreover, individuals with 4G alleles may respond more favorably to training in terms of improving fibrinolysis.

TI Effects of a common genetic polymorphism and exercise training on fibrinolysis in men and women aged 50--70

SO Dissertation Abstracts International, (2001) Vol. 62, No. 12B, p. 5635. Order No.: AAI3035829. 172 pages.

ISBN: 0-493-48931-2.

AB

Prospective epidemiological investigations suggest that impaired **fibrinolysis** is an independent risk factor for cardiovascular disease. Endurance **exercise** training has an inconsistent effect on **fibrinolysis**, possibly due to inadequate study design and lack of control of critical variables. The purpose of this study was to investigate the effects of endurance **exercise** training on blood **fibrinolysis** in middle-aged to older men and women.

The plasminogen activator inhibitor-1 (PAI-1) gene promoter 4G/5G polymorphism appears to affect circulating PAI-1 levels. The present study also determined if the PAI-1 polymorphism and initial PAI-1 levels play roles in **exercise** training-induced **fibrinolysis** changes.

Sedentary healthy volunteers aged 50-70 years were studied. Subjects completed a 6-week dietary program before baseline testing. Baseline tests in 48 subjects included fibrinolytic enzymes (tissue plasminogen activator (t-PA) and PAI-1 **activities**, and t-PA antigen), lipid-lipoprotein profiles, body composition, and maximal oxygen consumption (VO₂ max). Genotyping was performed using PCR/RFLP techniques. Baseline tests were repeated in 34 subjects after 6-month **exercise** intervention.

Baseline fibrinolytic levels were not significantly different among genotype groups (15.12 ± 1.39 vs. 11.97 ± 0.73 vs. 11.89 ± 1.39 AU/mL, for 4G/4G, 4G/5G, and 5G/5G, respectively). However, after controlling for triglyceride levels, the 4G/4G genotype had significantly higher PAI-1 **activity** levels than the other genotypes. After **exercise** training, **fibrinolysis** improved as shown by a 30% increase in t-PA **activity** (0.18 ± 0.07 IU/mL, $p = 0.016$) and a 6% decrease in t-PA antigen (-0.58 ± 0.20 , $p = 0.006$). Subgroup analyses revealed a significant 41% increase ($p = 0.03$) in t-PA **activity** levels in the 4G/5G group and a 10% decrease in t-PA antigen levels ($p = 0.008$) in the 4G/4G group. Training did not change PAI-1 **activity** levels significantly. ANOVAs revealed no differences in fibrinolytic measure changes with training among genotype groups. Adjusting for initial levels did not change results for PAI-1 **activity** and t-PA antigen. Interestingly, there was a significant interaction between baseline t-PA **activity** and genotype in determining final t-PA **activity**.

The present results suggest that endurance **exercise** training may increase **fibrinolysis** in middle-aged to older men and women by increasing t-PA **activity** and decreasing t-PA antigen levels. Moreover, individuals with 4G alleles may respond more favorably to training in terms of improving **fibrinolysis**.

Kapushoc, Stephen T.

From: Catlin, Karen (ASRC)
Sent: Wednesday, February 15, 2006 10:44 AM
To: Kapushoc, Stephen T.
Subject: FW: Dissertation availability

Hi again,

See below response from the electronic publisher of dissertations re: the availability date of Tiyasangthong's dissertation.

Please let me know if you need anything else.

Karen

-----Original Message-----

From: Heyboer, Tonya [mailto:Tonya.Heyboer@il.proquest.com]
Sent: Wednesday, February 15, 2006 9:22 AM
To: Catlin, Karen (ASRC)
Subject: RE: Dissertation availability

Hi Karen,

I found the dissertation you reference. The citation lists June 2002, which means this is the month we first published this dissertation in our DAI print index. Some at PTO have told me that is what they look for as the "publication date". But clearly this thesis was written and defended earlier than that. 2001 (which you see on the results list) was the degree year. So it looks like for whatever reason the author/institution chose to wait before sending to us. Unfortunately, that's as much information as we have.

Tonya

Tonya Heyboer
Trainer - Government, Corporate & Medical Accounts
ProQuest Information and Learning
Phone: 540-885-8688
Mobile: 540-230-0997
Email: tonya.heyboer@il.proquest.com

-----Original Message-----

From: Catlin, Karen (ASRC) [mailto:Karen.Catlin@USPTO.GOV]
Sent: Wednesday, February 15, 2006 9:00 AM
To: Heyboer, Tonya
Subject: Dissertation availability

Good morning Tonya,

I have a question about the date a specific dissertation was available via ProQuest Dissertations and Theses. Is it possible to determine the date that "Effects of a common genetic polymorphism and exercise training on fibrinolysis in men and women aged 50-70" was available? The ISBN is 0493489312, and the Order No. is AAI3035829. The only date I could find was the year 2001 which is on the document itself. The examiner requesting the information would a more specific date. The University of Maryland, where the dissertation was presented, has not replied to my question about an exact date.

Thanks for your help, and thanks again for the demos last week!
Karen

Karen Catlin (ASRC Aerospace)
Web Projects/Reference Librarian
Scientific and Technical Information Center
US Patent and Trademark Office
571-272-2573